

How Environmental Exposures Influence the Development and Exacerbation of Asthma

Ruth A. Etzel, MD, PhD

ABSTRACT. Environmental exposures may increase a child's risk of developing asthma and also may increase the risk of asthma exacerbations. This article reviews several environmental exposures and suggests whether they contribute to asthma prevalence, asthma exacerbations, or both. Outdoor air exposures and violence are not likely to cause the increase in asthma prevalence. Exposure to outdoor air pollutants primarily leads to increased exacerbations, sometimes manifested as asthma clusters. Clinicians should be alert for space-time clusters of asthma exacerbations in the community, because these clusters may suggest a modifiable point-source exposure. Indoor air exposures are more strongly linked to the increase in asthma prevalence. Exposure to dust mites and tobacco smoke are risk factors for the development of asthma and may also exacerbate existing asthma. Effective measures to prevent exposures to these pollutants are available. With proper management, the amount of environmental exposures can be decreased. Whether decreasing these exposures will result in decreases in asthma prevalence and exacerbations is not yet documented. *Pediatrics* 2003;112:233–239; *asthma, indoor air pollution, outdoor air pollution.*

ABBREVIATION. ppb, parts per billion.

Rates of asthma morbidity and mortality are increasing.^{1–7} From 1980 to 1994, the prevalence of asthma in the United States increased 75%.⁸ Asthma is more prevalent among black children than among white children.^{2,9} Black children are hospitalized for asthma at a higher rate than are white children, although much of this difference is likely attributable to poverty rather than to race.^{10,11}

From a public health perspective, it is useful to consider 2 separate dimensions of the asthma problem. The first is that the proportion of children in whom asthma has been diagnosed is increasing. Environmental exposures are among the many possibilities that have been proposed to explain the increase. The second component of the asthma problem is the increase in asthma attacks among children who already have asthma. Environmental exposures are also among the many possible explanations for this increase.

For the purposes of this article, the home environment includes exposures inside the home (eg, dust mites, cats and dogs, cockroaches, environmental tobacco smoke, molds) and also exposures in the neighborhood of the home (eg, outdoor air pollutants, violence). A child's home environment contributes to the risk of developing asthma and the subsequent risk of having asthma attacks^{12–14}; a recent study indicated that violence may also contribute to the risk of asthma attacks.¹⁵ Children who live in poverty are often exposed to violence. Whether children who live in poverty are more heavily exposed to indoor and outdoor air pollutants than are children who do not live in poverty is not known.

It is essential for clinicians to be knowledgeable about environmental precipitants of asthma, because this information may help them to counsel patients and their parents. Ideally, clinicians would focus their energy on primary prevention of asthma (prevention of the initial onset of asthma). At this time, however, the effectiveness of many primary prevention strategies is unknown. As a result, most clinicians have relied on secondary prevention (ie, trying to prevent exacerbations in children with known asthma, in part by decreasing environmental exposures, which cause worsening of symptoms). What is not widely recognized, however, is that the effectiveness of many secondary prevention measures is also unknown.

The purpose of this article is to review the evidence regarding the link between environmental exposures and asthma prevalence and exacerbations. The realization that environmental exposures could lead to asthma attacks has its origins in studies of clusters of asthma.

Much of what we now know about the relation between outdoor air exposures and asthma came to light as a result of asthma clusters in communities. Clinicians should be alert for space-time clusters of asthma exacerbations in the community, because these clusters may suggest a modifiable point-source exposure.^{16–19} Specific outdoor air pollutants linked to clusters of asthma are described in the following paragraphs.

EXPOSURE TO OUTDOOR AIR POLLUTANTS

Exposure to Castor Bean Dust

One of the first studies to document the relationship between air pollution and asthma was an investigation of an asthma cluster in Ohio. In the late 1920s, a cluster of asthma affecting 200 patients oc-

From the Department of Environmental and Occupational Health, School of Public Health and Health Services, George Washington University, Washington, DC.

Received for publication Jul 26, 2002; accepted Jan 17, 2003.

Address correspondence to Ruth A. Etzel, MD, PhD, 4320 Diplomacy Dr, Ste 2630, Anchorage, AK 99508. E-mail: retzel@earthlink.net

PEDIATRICS (ISSN 0031 4005). Copyright © 2003 by the American Academy of Pediatrics.

curred in Toledo.¹⁶ Most affected patients lived within a 1-mile radius of a castor bean mill, which also produced linseed oil. Many of the patients claimed that their attacks coincided with the odor of linseed oil coming from the mill when the wind blew in the right direction. When the health department investigated the possibility that linseed oil might be causing the asthma attacks, they considered it unlikely because the seed was too heavy to become windborne. The mill, however, not only manufactured linseed oil but also expressed castor oil from castor beans. The castor beans produced a fine dust that was readily carried by the wind into the surrounding neighborhoods. The outbreak of asthma attacks in Toledo was documented to have been caused by the inhalation of castor bean grinding dust, and the epidemic disappeared after the factory stopped processing castor beans.¹⁶ Asthma epidemics also have been documented to occur in other locations in the United States, including Brooklyn, New York,²⁰ but the cause of these epidemics is elusive.

Exposure to Grain Dust

Beginning in 1953 and continuing for nearly 20 years, Charity Hospital in New Orleans experienced asthma epidemics. These epidemics often involved >100 asthma-related visits to the emergency department in a single 24-hour period. In 1955, >350 patients with asthma were treated in the emergency department in a period of <24 hours; 2 patients died. In early November 1960, >200 people with asthma sought care in the emergency department in a single day. Epidemics such as this continued to occur until 1968, when they stopped, for reasons that were not fully understood. A large number of possible causes were studied.²¹⁻²⁵ One of the first hypotheses was that the asthma outbreaks might be attributable to inhalation of particles from a waste dump site where spontaneous underground burning was occurring.^{21,26} When the waste dump was removed, the asthma outbreaks continued.^{22,27} Other possibilities were suggested, including fire smoke emissions and aeroallergens.²⁸ Grain dust was also considered,²⁹ because a large amount of grain from Midwestern farms is shipped on barges down the Mississippi River and exported through New Orleans. Along the short stretch of the river between Baton Rouge and the Gulf of Mexico, there are >60 loading facilities. In 1968, the year the epidemics stopped, the grain elevators in New Orleans were modernized. The modernization included the addition of new dust control equipment, such as air filters on the silos. It is likely that these filters prevented soy dust from reaching the city's residents. An ecologic study documented that the asthma epidemics occurred on days when barges carrying soybeans were unloading in the New Orleans harbor.³⁰

Exposure to Soybean Dust

In the early 1980s, emergency department physicians in Barcelona, Spain, documented that sudden increases in visits for acute severe asthma overwhelmed the emergency services on certain days.

Between 1981 and 1987, 1155 emergency department visits by 687 people occurred on 26 asthma epidemic days. Careful daily surveillance was undertaken, with documentation of the place, day, and time when each person seeking emergency department care at each of 4 large hospitals experienced an asthma attack. Geographic mapping of the place where most emergency department visitors experienced attacks on epidemic days demonstrated that symptoms were most likely to start in the central part of the city, near the harbor. Epidemic asthma was found to occur only on weekdays, never on weekends. Furthermore, on epidemic days, most people experienced their attacks in the middle of the day, usually between 11 AM and 1 PM. After studying each of the products loaded or unloaded from ships in the harbor, it was determined that the outbreaks of asthma were caused by inhalation of soybean dust released from silos during the unloading of soybeans at the city harbor.^{17,18} In 1997, filters were placed on the silos, decreasing the amount of soybean dust emitted into the ambient air. No additional epidemics of asthma were reported in Barcelona.¹⁹

Exposure to Wood Smoke

Smoke from bush fires has been linked to increases in emergency department visits for asthma in Sydney, Australia,³¹ and smoke from forest fires has been linked to increases in emergency department visits for asthma in California.³² These and other studies indicate that localized outdoor air pollution may play an important role in asthma.

Exposure to Ambient Air Pollution

Ambient (outdoor) air pollution may be deleterious to the health of children with asthma.³³⁻⁴⁰ Some children with asthma (as well as some children without asthma) have decreases in lung function after exposure to ozone.⁴¹⁻⁴⁵ Ozone, a pollutant that is formed primarily by vehicular exhaust and is the principal component of urban smog, is associated with asthma exacerbations in some children with reactive airway disease. Levels of ozone are usually greatest on hot summer days, and the levels tend to reach their peak in the late afternoon. Exposure to ambient sulfur oxides and suspended particulates may also lead to pulmonary function decreases in children.⁴⁶⁻⁵²

Nitrogen dioxide is an oxidant gas that can penetrate deep into the lungs and damage delicate lung tissues. Some studies have shown a relationship between nitrogen dioxide and respiratory symptoms. Shima et al⁵³ have demonstrated that the prevalence of bronchitis, wheezing, and asthma increased with each increase of 10 parts per billion (ppb) in indoor nitrogen dioxide concentrations among girls but not among boys.

Epidemiologic studies have documented that the relationship between air pollution and hospital admissions for respiratory illnesses may be the largest among people in inner cities.⁵⁴ White et al⁵⁵ demonstrated that the average number of emergency department visits for asthma or reactive airways disease among inner-city children in Atlanta was 37%

greater on days when the maximum ozone levels exceeded 11 ppb. Tolbert et al⁵⁶ showed that the relative risk for a pediatric emergency department visit increased by 1.04 per increase of 20 ppb in the maximum 8-hour ozone exposure level.

Although ambient air pollution may exacerbate asthma among individual children, a large international study suggests that outdoor air pollution is not a major factor in the development of asthma in populations.⁵⁷ It is not likely that outdoor air pollution could account for the increasing asthma prevalence in children.

EXPOSURE TO INDOOR AIR POLLUTION

Exposure to indoor air pollutants may have a more important effect on the development of childhood asthma than may exposure to outdoor air pollutants.⁵⁸ Many allergens and irritants (from smoke, cockroaches, mites, molds, cats, and dogs) are found indoors.⁵⁸⁻⁶¹ In the United States, most children spend the majority of each day (average 20 hours) inside buildings.⁶² Indoor air pollution has become a particular problem since the energy crisis of the 1970s, which led to the construction of more energy-efficient buildings with less air circulation.

The Institute of Medicine recently released a report, "Clearing the Air," on the relationship between indoor air pollution and asthma.⁵⁸ The report described the quality of the evidence supporting the relationship between certain indoor pollutants and asthma development and exacerbations. Five levels of evidence were identified: sufficient evidence of a causal relationship, sufficient evidence of an association, limited or suggestive evidence of an association, inadequate or insufficient evidence to determine whether an association exists, and limited or suggestive evidence of no association. Indoor air pollutants for which there is sufficient evidence of a causal relationship and sufficient evidence of an association are described in this article and are summarized in Table 1.

Exposure to Environmental Tobacco Smoke

Exposure to environmental tobacco smoke is a risk factor for the development of asthma and for asthma attacks in children with existing disease. The Institute of Medicine concluded that there is sufficient evidence of an association between exposure to environmental tobacco smoke and the development of asthma and sufficient evidence of a causal relation-

ship between exposure to environmental tobacco smoke and exacerbations of asthma.⁵⁸

Birth certificate data for 1999 indicate that 12.3% of women who gave birth reported smoking during pregnancy.⁶³ Exposure to tobacco smoke products in utero is a risk factor for wheezing in the first year of life.⁶⁴

Approximately 42% of children 2 months to 11 years of age live in a home with at least 1 smoker.⁶⁵ An estimated 8.7 to 12.4 million US children younger than 5 years are exposed to cigarette smoke at home.⁶⁶ Children who have asthma and whose parents smoke have more frequent asthma attacks and more severe symptoms.⁶⁷⁻⁸⁰ One study suggested that if parents expose their children with asthma to less cigarette smoke, then their asthma symptoms may be less severe.⁸¹ More studies are needed on the effectiveness of decreasing the amount of environmental tobacco smoke to which children are exposed. Nonetheless, the Centers for Disease Control and Prevention has conducted systematic reviews on 11 interventions to decrease the amount of environmental tobacco smoke to which children are exposed and has made recommendations regarding the use of these interventions.⁸²

Indoor Exposure to Dust Mites

The Institute of Medicine has concluded that sensitization to house dust mites is an important risk factor for asthma development and also asthma exacerbations.^{58,83-85} Household interventions can decrease children's exposure to dust mites. Plastic mattress covers are an effective measure to decrease dust mite infestation of bedding. Several randomized controlled trials have demonstrated a significant decrease in concentrations of mite allergen on mattresses covered by polyurethane casings.^{86,87}

Vojta et al⁸⁸ recently conducted a randomized trial of physical interventions to decrease house dust mite allergen levels in low-income urban homes. They studied bedroom carpet interventions in 11 homes. Six homes received intensive vacuuming only, and 5 homes received dry steam cleaning and intensive vacuuming. Although both groups of homes had significant decreases in house dust mite levels in the bedroom carpets, only the homes that had received dry steam cleaning had decreases in house dust mite levels that persisted for up to 8 weeks after the intervention. They also studied bed interventions in 11 homes. All 11 beds received allergen-impermeable

TABLE 1. Relationship Between Indoor Exposures and Asthma Development and Exacerbations

Indoor Exposure	Asthma Development	Asthma Exacerbations
House dust mite	Increased	Increased
Environmental tobacco smoke	Probably increased	Increased
Cockroach	Maybe increased	Increased
Cat	Maybe increased	Increased
Dog	Maybe increased	Probably increased
Molds	?	Probably increased
Oxides of nitrogen	?	Probably increased
Ozone	?	Probably increased
Particulates	?	Probably increased
Sulfur dioxide	?	Probably increased

box springs, mattresses, and pillow covers. The bedding in 6 homes was laundered weekly by a professional service, and the bedding in the other 5 homes was laundered at home. Both interventions resulted in significant decreases in house dust mite levels in the beds. They also evaluated the effects of dry steam cleaning versus intensive vacuuming of upholstered furniture. Both interventions were about equally effective in decreasing dust mite levels in upholstered furniture.

Other investigators have shown little or no impact of house dust mites on asthma. Lau et al⁸⁹ showed no correlation between levels of house dust mites and asthma in children at 7 years of age. Gotzsche et al⁹⁰ performed a meta-analysis of 23 studies of house dust mite control measures in the management of childhood and adult asthma. Five of the 23 studies showed a decrease in dust mite exposure. When all studies were considered, the results did not show any clinical benefit from measures to reduce dust mites.^{90,91} The same was true when only the 5 trials that showed a decrease in dust mite exposure were considered. Additional studies of effectiveness of dust mite reduction techniques in homes of children with asthma are needed.

Exposure to Cockroaches

The Institute of Medicine concluded that cockroach allergens are causally related to asthma attacks.⁵⁸ Cockroach droppings may be one of the most underappreciated allergens in the indoor environment.⁹² Multicenter asthma studies funded by the National Institutes of Health have brought to light the importance of cockroach allergens in causing morbidity among inner-city children with asthma. Cockroach allergens not only increase the risk of asthma attacks but also may increase a child's risk of developing asthma.^{58,93} In a study among 235 children in Boston, children who lived with higher levels of cockroach antigen (as measured by *Bla g* levels from kitchen samples) were 4 times more likely to have a new diagnosis of asthma than were children from homes with a low level of cockroach antigen.⁹³

Exposure to Cats

The Institute of Medicine concluded that exposure to cats is causally related to asthma exacerbations among children with asthma.⁵⁸ Recent studies,^{94,95} however, have shown that the presence of a cat in the house may decrease the risk of developing asthma. The reasons for this apparently contradictory finding are being investigated.

Exposure to Molds

Indoor Exposure

The Institute of Medicine concluded that exposure to molds is associated with asthma exacerbations.⁵⁸ Some molds may invoke an allergic response (resulting in asthma or allergic rhinitis) in susceptible children. Exposure to molds may lead to allergic sensitization.⁶¹ At least 60 species of molds have spores thought to be allergenic.⁹⁶ Species of particular concern are *Penicillium*, *Aspergillus*, *Cladosporium*, and

Alternaria. On exposure to these species, nasal congestion, runny nose, sneezing, conjunctivitis, lacrimation, wheezing, chest tightness, and shortness of breath may occur. Thirty percent of patients with respiratory allergies seem to be particularly sensitive to molds. Children are the most sensitive population.⁹⁷⁻⁹⁹ Strong associations have been found between mold or dampness and respiratory symptoms (eg, wheezing, sore throat, runny nose)^{97,98,100-105} and doctor-diagnosed asthma.¹⁰⁶⁻¹⁰⁸

Outdoor Exposure

Fungal concentrations in outdoor air vary according to the season and weather conditions.¹⁰⁹ In subarctic climates, patients with mold allergies have more serious symptoms during spring and autumn when outdoor mold concentrations are usually highest.¹¹⁰ The 1997 National Heart, Lung, and Blood Institute *Guidelines for the Diagnosis and Management of Asthma* suggest that clinicians who care for children with asthma inquire about molds in the home environment and suggest methods to decrease the amount of mold to which children are exposed.¹¹¹

Exposure to environmental molds has been documented to play a role in asthma-related mortality.¹¹² Airborne molds have been linked to asthma deaths among 5- to 34-year-olds in Chicago.¹¹³ The odds of an asthma death occurring on days with mold spore counts of ≥ 1000 spores/m³ was 2.16 times higher (95% confidence interval: 1.31-3.56) than that on days with mold spore counts <1000 spores/m³.

Exposure to Violence

A potential risk factor for asthma that has recently been recognized is violence. Stress may potentiate the allergic response to allergens by increasing the release of inflammatory mediators and the subsequent cascade of inflammatory events characteristic of chronic asthma.^{15,114}

PROTECTIVE FACTORS

Several risk factors have been identified as protective against asthma. Ball et al¹⁴ showed that exposure of young children to older children at home or to other children in child care settings protects against the development of asthma and frequent wheezing later in childhood. They hypothesized that within the first 6 months of an infant's life, the immune response of children without atopy shifts from one associated predominantly with type 2 helper T cells, such as that in adults with atopic illnesses, toward one based more on cytokines derived from type 1 helper T cells, such as that in adults without atopy.^{14,115} A recent study has shown that long-term and early-life exposure to stables and farm milk induces a strong protective effect against development of asthma, hay fever, and atopic sensitization.¹¹⁶

PREVENTION

The aforementioned studies have documented that specific interventions can decrease environmental exposures in homes of children with asthma. Despite the growing body of evidence, many children and their families, particularly children who live in pov-

erty and rely on emergency departments as their primary source of health care, may not be receiving adequate counseling about how to avoid environmental exposures. A variety of excellent educational materials have been developed to teach parents and children with asthma about the importance of their environment, and several excellent reviews of indoor environmental controls in asthma management have been published.^{117–119}

To prevent unnecessary exposures to outdoor air pollution, clinicians who care for children should be aware of the implications of municipal smog alerts and provide appropriate guidance to children with asthma and their parents regarding exercise during periods of high pollution.^{120,121}

With proper management, many environmental exposures can be decreased. What has yet to be documented is whether this will result in decreased morbidity from asthma. Studies to investigate this hypothesis are currently under way.

REFERENCES

- Evans R III, Mullally DI, Wilson RW, et al. National trends in the morbidity and mortality of asthma in the US. Prevalence, hospitalization, and death from asthma over two decades: 1965–1984. *Chest*. 1987;91(suppl):655–745
- Gergen PJ, Mullally DI, Evans R III. National survey of prevalence of asthma among children in the United States, 1976 to 1980. *Pediatrics*. 1988;81:1–7
- Sly RM. Mortality from asthma, 1979–1984. *J Allergy Clin Immunol*. 1988;82:705–717
- Centers for Disease Control and Prevention. Asthma—United States, 1980–1987. *MMWR Morb Mortal Wkly Rep*. 1990;39:493–497
- Brown CM, Anderson HA, Etzel RA. Asthma. The states' challenge. *Public Health Rep*. 1997;112:198–205
- Vollmer WM, Osborne ML, Buist AS. 20-year trends in the prevalence of asthma and chronic airflow obstruction in an HMO. *Am J Respir Crit Care Med*. 1998;157:1079–1084
- Holgate ST. The epidemic of allergy and asthma. *Nature*. 1999;402(6760 suppl):B2–B4
- Centers for Disease Control and Prevention. Surveillance for asthma—United States, 1960–1995. *CDC Surveill Summ*. 1998;47:1–28
- Weitzman M, Gortmaker SL, Sobol AM, Perrin JM. Recent trends in the prevalence and severity of childhood asthma. *JAMA*. 1992;268:2673–2677
- Gergen PJ, Weiss KB. Changing patterns of asthma hospitalization among children: 1979 to 1987. *JAMA*. 1990;264:1688–1692
- Wissow LS, Gittelsohn AM, Szklo M, Starfield B, Mussman M. Poverty, race, and hospitalization for childhood asthma. *Am J Public Health*. 1988;78:777–782
- Cullinan P, Taylor AJ. Asthma in children: environmental factors. *BMJ*. 1994;308:1585–1586
- Holt PG, Macaubas C, Stumbles PA, Sly PD. The role of allergy in the development of asthma. *Nature*. 1999;42(6760 suppl):B12–B17
- Ball TM, Castro-Rodriguez JA, Griffith KA, Holberg CJ, Martinez FD, Wright AL. Siblings, day-care attendance, and the risk of asthma and wheezing during childhood. *N Engl J Med*. 2000;343:538–543
- Wright RJ, Steinbach SF. Violence: an unrecognized environmental exposure that may contribute to greater asthma morbidity in high risk inner-city populations. *Environ Health Perspect*. 2001;109:1085–1089
- Figley KD, Elrod RH. Endemic asthma due to castor bean dust. *JAMA*. 1928;90:79–82
- Sunyer J, Anto JM, Rodrigo MJ, Morrell F. Case-control study of serum immunoglobulin-E antibodies reactive with soybean in epidemic asthma. *Lancet*. 1989;1:179–182
- Anto JM, Sunyer J, Rodriguez-Roisin R, Suarez-Cervera M, Vazquez L. Community outbreaks of asthma associated with inhalation of soybean dust. *N Engl J Med*. 1989;320:1097–1102
- Anto JM, Sunyer J, Reed CE, et al. Preventing asthma epidemics due to soybeans by dust-control measures. *N Engl J Med*. 1993;329:1760–1763
- Goldstein IF, Currie B. Seasonal patterns of asthma: a clue to etiology. *Environ Res*. 1984;33:201–215
- Lewis R, Gilkeso M, McCamdin RO. Air pollution and New Orleans asthma. *Public Health Rep*. 1962;77:947–954
- Carroll RE. Environmental epidemiology. V. Epidemiology of New Orleans epidemic asthma. *Am J Public Health Nations Health*. 1968;58:1677–1683
- Booth S, DeGroot I, Markush R, Horton RJ. Detection of asthma epidemics in seven cities. *Arch Environ Health*. 1965;10:152–155
- Salvaggio J, Hasselblad V, Seabury J, Heiderscheid LT. New Orleans asthma. II. Relationship of climatologic and seasonal factors to outbreaks. *J Allergy*. 1970;45:257–265
- Goldstein IF. Weather patterns and asthma epidemics in New York City and New Orleans, USA. *Int J Biometeorol*. 1980;24:329–339
- Weill H, Ziskind MM, Dickerson RC, Derbes VJ. Epidemic asthma in New Orleans. *JAMA*. 1964;190:811–814
- Salvaggio JE, Klein RC. New Orleans asthma. I. Characterization of individuals involved in epidemics. *J Allergy*. 1967;39:227–233
- Hendrick DJ. Asthma: epidemics and epidemiology. *Thorax*. 1989;44:609–613
- Weill H, Ziskind MM, Dickerson RC, Derbes VJ. Allergic air pollutants in New Orleans. *J Air Pollut Control Assoc*. 1965;15:467–471
- White MC, Etzel RA, Olson DR, Goldstein IF. Reexamination of epidemic asthma in New Orleans, Louisiana, in relation to the presence of soy at the harbor. *Am J Epidemiol*. 1997;145:432–438
- Churches T, Corbett S. Asthma and air pollution in Sydney. *N S W Public Health Bull*. 1991;2:72–73
- Duclos P, Sanderson LM, Lipsett M. The 1987 forest fire disaster in California: assessment of emergency room visits. *Arch Environ Health*. 1990;45:53–58
- Bates DV, Sizto R. Air pollution and hospital admissions in southern Ontario, the acid summer haze effect. *Environ Res*. 1987;43:317–331
- Bates DV. The effects of air pollution on children. *Environ Health Perspect*. 1995;103:49–53
- Burnett RT, Smith-Doiron M, Raizenne ME, Stieb D. Association between ozone and hospitalization for acute respiratory diseases in children less than 2 years of age. *Am J Epidemiol*. 2001;153:444–452
- American Academy of Pediatrics, Committee on Environmental Health. Ambient air pollution: respiratory hazards to children. *Pediatrics*. 1993;91:1210–1213
- Committee of the Environmental and Occupational Health Assembly of the American Thoracic Society. Health effects of outdoor air pollution. Part 2. *Am J Respir Crit Care Med*. 1996;153:477–498
- Committee of the Environmental and Occupational Health Assembly of the American Thoracic Society. Health effects of outdoor air pollution. *Am J Respir Crit Care Med*. 1996;153:3–50
- Ostro B, Lipsett M, Mann J, Braxton-Owens H, White M. Air pollution and exacerbation of asthma in African-American children in Los Angeles. *Epidemiology*. 2001;12:200–208
- McConnell R, Berhane K, Gilliland F, et al. Air pollution and bronchitic symptoms in Southern California children with asthma. *Environ Health Perspect*. 1999;107:757–760
- Avol EL, Linn WS, Shamoo DA, et al. Respiratory effects of photochemical oxidant air pollution in exercising adolescents. *Am Rev Respir Dis*. 1985;132:619–622
- Avol EL, Linn WS, Shamoo DA, et al. Short-term respiratory effects of photochemical oxidant exposure in exercising children. *J Air Pollut Control Assoc*. 1987;37:158–162
- McDonnell WF III, Chapman RS, Leigh MW, Strobe GL, Collier AM. Respiratory responses of vigorously exercising children to 0.12 ppm ozone exposure. *Am Rev Respir Dis*. 1985;132:875–879
- Spektor DM, Lippman M, Liou PJ, et al. Effects of ambient ozone on respiratory function in active, normal children. *Am Rev Respir Dis*. 1988;137:313–320
- Spektor DM, Thurston GD, Mao J, He D, Hayes C, Lippmann M. Effects of single- and multiday ozone exposures on respiratory function in active normal children. *Environ Res*. 1991;55:107–122
- Ware JH, Ferris BG Jr, Dockery DW, Spengler JD, Stram DO, Speizer FE. Effects of ambient sulfur oxides and suspended particles on respiratory health of preadolescent children. *Am Rev Respir Dis*. 1986;133:834–842
- Dassen W, Brunekreef B, Hoek G, et al. Decline in children's pulmonary function during an air pollution episode. *J Air Pollut Control Assoc*. 1986;36:1223–1227
- Dockery DW, Speizer FE, Stram DO, Ware JH, Spengler JD, Ferris BG Jr. Effects of inhalable particles on expiratory health of children. *Am Rev Respir Dis*. 1989;139:587–594
- Koenig JQ, Covert DS, Pierson WE. Effects of inhalation of acidic compounds on pulmonary function in allergic adolescent subjects. *Environ Health Perspect*. 1989;79:173–178
- Koenig JQ, Pierson WE, Horike M. The effects of inhaled sulfuric acid

- on pulmonary function in adolescent asthmatics. *Am Rev Respir Dis.* 1983;128:221-225
51. Koenig JQ, Covert DS, Hanley QS, et al. Prior exposure to ozone potentiates subsequent response to sulfur dioxide in adolescent asthmatic subjects. *Am Rev Respir Dis.* 1990;141:377-380
 52. Raizenne ME, Burnett RT, Stern B, Franklin CA, Spengler JD. Acute lung function responses to ambient acid aerosol exposures in children. *Environ Health Perspect.* 1989;79:179-185
 53. Shima M, Adachi M. Effect of outdoor and indoor nitrogen dioxide on respiratory symptoms in schoolchildren. *Int J Epidemiol.* 2000;29:862-870
 54. Thurston GD, Ito K, Kinney PL, Lippmann M. A multi-year study of air pollution and respiratory hospital admissions in three New York State metropolitan areas: results for the 1988 and 1989 summers. *J Expo Anal Environ Epidemiol.* 1992;2:429-450
 55. White MC, Etzel RA, Wilcox WD, Lloyd C. Exacerbations of childhood asthma and ozone pollution in Atlanta. *Environ Res.* 1994;65:56-68
 56. Tolbert PE, Mulholland JA, MacIntosh DL, et al. Air quality and pediatric emergency room visits for asthma in Atlanta, Georgia, USA. *Am J Epidemiol.* 2000;151:798-810
 57. International Study of Asthma and Allergies in Childhood Steering Committee. Worldwide variation in prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and atopic eczema: ISAAC. *Lancet.* 1998;351:1225-1232
 58. Institute of Medicine, Committee on the Assessment of Asthma and Indoor Air. *Clearing the Air: Asthma and Indoor Air Exposures.* Washington, DC: National Academy Press; 2000
 59. Etzel RA. Indoor air pollution and childhood asthma: effective environmental interventions. *Environ Health Perspect.* 1995;103(suppl 6):55-58
 60. Platts-Mills TA. Allergen-specific treatment for asthma: III. *Am Rev Respir Dis.* 1993;148:553-555
 61. Pope AM, Patterson R, Burge H, et al, eds. *Indoor Allergens: Assessing and Controlling Adverse Health Effects.* Institute of Medicine, Committee on the Health Effects of Indoor Allergens. Washington, DC: National Academy Press; 1993
 62. Schwab M, McDermott A, Spengler JD. Using longitudinal data to understand children's activity patterns in an exposure context: Data from the Kanawha County Health Study. *Environ Int.* 1992;18:173-189
 63. Matthews TJ. Smoking during pregnancy in the 1990s. *Natl Vital Stat Rep.* 2001;49:1-14
 64. Tager IB, Hanrahan JP, Tosteson TD, et al. Lung function, pre- and post-natal smoke exposure, and wheezing in the first year of life. *Am Rev Respir Dis.* 1993;147:811-817
 65. Pirkle JL, Flegal KM, Bernert JT, Brady DJ, Etzel RA, Maurer KR. Exposure of the U.S. population to environmental tobacco smoke: the Third National Health and Nutrition Examination Survey, 1988-1991. *JAMA.* 1996;275:1233-1240
 66. American Academy of Pediatrics, Committee on Environmental Hazards. Involuntary smoking—a hazard to children. *Pediatrics.* 1986;77:755-757
 67. Murray AB, Morrison BJ. The effect of cigarette smoke from the mother on bronchial responsiveness and severity of symptoms in children with asthma. *J Allergy Clin Immunol.* 1986;77:575-581
 68. Evans D, Levison J, Feldman CH, et al. The impact of passive smoking on emergency room visits of urban children with asthma. *Am Rev Respir Dis.* 1987;135:567-572
 69. Burchfiel CM, Higgins MW, Keller JB, Howatt WF, Butler WJ, Higgins IT. Passive smoking in childhood: respiratory conditions and pulmonary function in Tecumseh, Michigan. *Am Rev Respir Dis.* 1986;133:966-973
 70. Chilmonczyk BA, Salmun LM, Megathlin KN, et al. Association between exposure to environmental tobacco smoke and exacerbations of asthma in children. *N Engl J Med.* 1993;328:1665-1669
 71. Ehrlich R, Kattan M, Godbold J, et al. Childhood asthma and passive smoking. Urinary cotinine as a biomarker of exposure. *Am Rev Respir Dis.* 1992;145:594-599
 72. Holberg CJ, Wright AL, Martinez FD, Morgan WJ, Taussig LM. Child day care, smoking by caregivers, and lower respiratory tract illness in the first 3 years of life. Group Health Medical Associates. *Pediatrics.* 1993;91:885-892
 73. Krzyzanowski M, Quackenboss JJ, Lebowitz MD. Chronic respiratory effects of indoor formaldehyde exposure. *Environ Res.* 1990;52:117-125
 74. Martinez FD, Cline M, Burrows B. Increased incidence of asthma in children of smoking mothers. *Pediatrics.* 1992;89:21-26
 75. Murray AB, Morrison BJ. Passive smoking by asthmatics: its greater effect on boys than girls and on older than on younger children. *Pediatrics.* 1989;84:451-459
 76. O'Connor GT, Weiss ST, Tager IB, Speizer FE. The effect of passive smoking on pulmonary function and nonspecific bronchial responsiveness in a population-based sample of children and young adults. *Am Rev Respir Dis.* 1987;135:800-804
 77. Oldigs M, Jorres R, Magnussen H. Acute effects of passive smoking on lung function and airway responsiveness in asthmatic children. *Pediatr Pulmonol.* 1991;10:123-131
 78. Rylander E, Pershagen G, Eriksson M, Bermann G. Parental smoking, urinary cotinine, and wheezing bronchitis in children. *Epidemiology.* 1995;6:289-293
 79. Sherman CB, Tosteson TD, Tager IB, Speizer FE, Weiss ST. Early childhood predictors of asthma. *Am J Epidemiol.* 1990;132:83-95
 80. Weitzman M, Gortmaker S, Walker DK, Sobol A. Maternal smoking and childhood asthma. *Pediatrics.* 1990;85:505-511
 81. Murray AB, Morrison BJ. The decrease in severity of asthma in children of parents who smoke since the parents have been exposing them to less cigarette smoke. *J Allergy Clin Immunol.* 1993;91:102-110
 82. Centers for Disease Control and Prevention. Strategies for reducing exposure to environmental tobacco smoke, increasing tobacco-use cessation, and reducing initiation in communities and health-care systems. *MMWR Morb Mortal Wkly Rep.* 2000;49(RR-12):1
 83. Sporik R, Holgate ST, Platts-Mills TA, Cogswell JJ. Exposure to house-dust mite allergen (Der p I) and the development of asthma in childhood. A prospective study. *N Engl J Med.* 1990;323:502-507
 84. Platts-Mills TA, Sporik RB, Wheatly LM, Heymann PW. Is there a dose-response relationship between exposure to indoor allergens and symptoms of asthma? *J Allergy Clin Immunol.* 1995;96:435-440
 85. Duffy DL, Mitchell CA, Martin NG. Genetic and environmental risk factors for asthma: a cotwin-control study. *Am J Respir Crit Care Med.* 1998;157:840-845
 86. Ehmert B, Lau-Schadendorf S, Weber A, Buettner P, Schou C, Wahn U. Reducing domestic exposure to dust mite allergen reduces bronchial hyperreactivity in sensitive children with asthma. *J Allergy Clin Immunol.* 1992;90:135-138
 87. Murray AB, Ferguson AC. Dust-free bedrooms in the treatment of asthmatic children with house dust or house dust mite allergy: a controlled trial. *Pediatrics.* 1983;71:418-422
 88. Vojta PJ, Randels SP, Stout J, et al. Effects of physical interventions on house dust mite allergen levels in carpet, bed, and upholstery dust in low-income, urban homes. *Environ Health Perspect.* 2001;109:815-819
 89. Lau S, Illi S, Sommerfeld C, et al. Early exposure to house-dust mite and cat allergens and development of childhood asthma: a cohort study. Multicentre Allergy Study Group. *Lancet.* 2000;356:1392-1397
 90. Gotzsche PC, Hammarquist C, Burr M. House dust mite control measures in the management of asthma: meta-analysis. *BMJ.* 1998;317:1105-1110
 91. Gotzsche PC, Johanson HK, Burr ML, Hammerquist C. House dust mite control measures for asthma. *Cochrane Database Syst Rev.* 2001;2:CD001187
 92. Rosenstreich D, Eggleston P, Katta M, et al. The role of cockroach allergy and exposure to cockroach allergen in causing morbidity among inner-city children with asthma. *N Engl J Med.* 1997;336:1356-1363
 93. Litonjua A, Weiss ST, Burge HA, Gold DR. Cockroach allergen levels and the incidence of asthma among children under five years old. *Am J Respir Crit Care Med.* 1998;157:A643
 94. Platts-Mills T, Vaughan J, Squillace S, Woodfolk J, Sporik R. Sensitization, asthma, and a modified Th2 response in children exposed to cat allergen: a population-based cross-sectional study. *Lancet.* 2001;357:752-756
 95. Nafstad P, Magnus P, Gaarder PI, Jaakkola JJ. Exposure to pets and atopy-related diseases in the first 4 years of life. *Allergy.* 2001;56:307-312
 96. Burge HA. Airborne allergenic fungi. Classification, nomenclature, and distribution. *Immunol Allergy Clin North Am.* 1989;9:307-319
 97. Martin CJ, Platt SD, Hunt SM. Housing conditions and ill health. *Br Med J (Clin Res Ed).* 1987;294:1125-1127
 98. Platt SD, Martin CJ, Hunt SJ, et al. Damp housing, mould growth, and symptomatic health state. *BMJ.* 1989;298:1673-1678
 99. Wanner H, Verhoeff A, Columbi A, et al. *Biological Particles in Indoor Environments.* Luxembourg City, Luxembourg: Office for Official Publications of the European Communities; 1993
 100. Dales RE, Zwanenburg H, Burnett R, Franklin CA. Respiratory health effects of home dampness and molds among Canadian children. *Am J Epidemiol.* 1991;134:196-203
 101. Strachan DP, Sanders CH. Damp housing and childhood asthma: respiratory effects of indoor air temperature and relative humidity. *J Epidemiol Community Health.* 1989;43:7-14

102. Strachan DP, Elton RA. Relationship between respiratory morbidity in children and the home environment. *Fam Pract.* 1986;3:137–142
103. Strachan DP. Damp housing and childhood asthma: validation of reporting symptoms. *BMJ.* 1988;297:1223–1226
104. Delfino RJ, Zeiger RS, Seltzer JM, et al. The effect of outdoor fungal spore concentrations on daily asthma severity. *Environ Health Perspect.* 1997;105:622–635
105. Delfino RJ, Coate BD, Zeiger RS, Seltzer JM, Street DH, Koutrakis P. Daily asthma severity in relation to personal ozone exposure and outdoor fungal spores. *Am J Respir Crit Care Med.* 1996;154:633–641
106. Husman T. Health effects of indoor-air microorganisms. *Scand J Work Environ Health.* 1996;22:5–13
107. Brunekreef B, Dockery DW, Speizer FE, Ware JH, Spengler JD, Ferris BG. Home dampness and respiratory morbidity in children. *Am Rev Respir Dis.* 1989;140:1363–1367
108. Dekker C, Dales R, Bartlett S, Brunekreef B, Zwanenburg H. Childhood asthma and the home environment. *Chest.* 1991;100:922–926
109. Hjelmroos M. Relationship between airborne fungal spore presence and weather variables: *Cladosporium* and *Alternaria*. *Grana.* 1993;32:40–47
110. Goldfarb AA. The significance of mold spore allergy in childhood respiratory disease. *Ann Allergy.* 1968;26:321–327
111. National Heart, Lung, and Blood Institute. *Guidelines for the Diagnosis and Management of Asthma.* Rockville, MD: National Institutes of Health; 1997 (NIH Publication No. 97-4051)
112. O'Hollaren MT, Yunginger JW, Offord KP, et al. Exposure to aeroallergens as a possible precipitating factor in respiratory arrest in young patients with asthma. *N Engl J Med.* 1991;324:359–363
113. Targonski PV, Persky VW, Ramekrishnan V. Effect of environmental molds on risk of death from asthma during the pollen season. *J Allergy Clin Immunol.* 1995;95:955–961
114. Wright RJ, Rodriguez M, Cohen S. Review of psychosocial stress and asthma: an integrated biopsychosocial approach. *Thorax.* 1998;53:1066–1074
115. Prescott SL, Macaubas C, Smallcombe T, Holt BJ, Sly PD, Holt PG. Development of allergen-specific T-cell memory in atopic and normal children. *Lancet.* 1999;353:196–200
116. Riedler J, Braun-Fahrlander C, Eder W, et al. Exposure to farming in early life and development of asthma and allergy: a cross-sectional survey. *Lancet.* 2001;358:1129–1133
117. Jones AP. Asthma and the home environment. *J Asthma.* 2000;37:103–124
118. Environmental controls and lung disease. Report of the ATS Workshop on Environmental Controls and Lung Disease. Santa Fe, New Mexico, March 24–25, 1988. *Am Rev Respir Dis.* 1990;142:915–939
119. Ingram JM, Heymann PW. Environmental controls in the management of asthma. *Immunol Allergy Clin North Am.* 1993;13:785–801
120. Etzel RA, Balk SJ, eds. American Academy of Pediatrics, Committee on Environmental Health. *Handbook of Pediatric Environmental Health.* Elk Grove Village, IL: American Academy of Pediatrics; 1999
121. Teague WC, Bayer CW. Outdoor air pollution. Asthma and other concerns. *Pediatr Clin North Am.* 2001;48:1167–1183

